

# LATTICE EXTENSIONS OF HECKE ALGEBRAS

IVAN MARIN

ABSTRACT. We investigate the extensions of the Hecke algebras of finite (complex) reflection groups by lattices of reflection subgroups that we introduced, for some of them, in our previous work on the Yokonuma-Hecke algebras and their connections with Artin groups. When the Hecke algebra is attached to the symmetric group, and the lattice contains all reflection subgroups, then these algebras are the diagram algebras of braids and ties of Aicardi and Juyumaya. We prove a structure theorem for these algebras, generalizing a result of Espinoza and Ryom-Hansen from the case of the symmetric group to the general case. We prove that these algebras are symmetric algebras at least when  $W$  is a Coxeter group, and in general under the trace conjecture of Broué, Malle and Michel.

## CONTENTS

1. Introduction	1
2. Structure	3
2.1. Semidirect extensions of group algebras by abelian algebras	3
2.2. Braid groups of reflection subgroups	4
2.3. Proof of the structure theorem	5
3. Traces	7
3.1. Reminder on canonical traces	7
3.2. Traces on generalized Hecke algebras	8
3.3. Symmetrizing trace	8
4. Main examples	8
4.1. The category of admissible semi-lattices and maps	9
4.2. The semi-lattice $\mathcal{L}_2$	9
4.3. The case of finite Coxeter groups	9
4.4. The parabolic lattice	10
4.5. Root systems	10
4.6. A priori unrelated examples	11
References	11

## 1. INTRODUCTION

Let  $W$  be finite complex reflection group, for instance a finite Coxeter group. Let  $B$  denote the braid group associated to  $W$  in the sense of Broué-Malle-Rouquier (see [10]), which in the case of a finite Coxeter group coincides with the Artin group attached to it. We denote  $\pi : B \rightarrow W$  the natural projection.

The object of this paper is to introduce and analyse a family of algebras denoted  $\mathcal{C}(W, \mathcal{L})$ , where  $\mathcal{L}$  is a finite join semi-lattice which lies inside the poset made of the full reflection subgroups of  $W$ , ordered by inclusion. Here a reflection subgroup of  $W$  is called *full* if, for any reflection in this subgroup, all the (pseudo-)reflections with the same reflecting hyperplane belong to it. The semi-lattice  $\mathcal{L}$  is additionally supposed to be stable under the natural action of  $W$  on the lattice of reflection subgroups, and to contain all the cyclic (full) reflection subgroups, and the trivial subgroup as well. Such a semi-lattice will be called an *admissible* semi-lattice.

---

*Date:* February 4, 2017.

*2010 Mathematics Subject Classification.* 20C08, 20F36, 20F55.

Let  $\mathcal{A}$  denote the hyperplane arrangement attached to  $W$ , namely the collection of its reflecting hyperplanes. Let  $\mathbf{k}$  be a commutative ring with 1, containing elements  $a_{H,i}$  where  $H \in \mathcal{A}$ ,  $0 \leq i < m_H$  where  $m_H$  is the order of the cyclic subgroup of  $W$  fixing  $H$ , with the convention that  $a_{H,i} = a_{w(H),i}$  for every  $H \in \mathcal{A}, w \in W$  and  $a_{H,0}$  is invertible inside  $\mathbf{k}$ . Let  $R$  denote the generic ring of Laurent polynomials with integer coefficients  $\mathbf{Z}[a_{H,i}, a_{H,0}^{\pm 1}]$ , with the same conventions. Our conditions on  $\mathbf{k}$  mean that it is a  $R$ -algebra. We now define  $\mathbf{k}$ -algebras  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$ , with the convention that  $\mathcal{C}(W, \mathcal{L}) = \mathcal{C}_R(W, \mathcal{L})$ .

These algebras are defined as follows. First consider the algebra  $\mathbf{k}\mathcal{L}$  defined as the free  $\mathbf{k}$ -module with basis elements  $e_\lambda, \lambda \in \mathcal{L}$ , and where the multiplication is defined by  $e_\lambda e_\mu = e_{\lambda \vee \mu}$ . This is sometimes called the Möbius algebra of  $\mathcal{L}$ . Elements of  $\mathcal{L}$  can be identified with the collection of reflecting hyperplanes attached to them, and we let  $e_H = e_{\{H\}}$  denote the idempotent attached to the subgroup fixing  $H \in \mathcal{A}$ . We shall use this identification whenever it is convenient to us.

By definition  $W$  acts by automorphisms on  $\mathbf{k}\mathcal{L}$ , hence so does  $B$ , and one can form the semidirect product  $\mathbf{k}B \ltimes \mathbf{k}\mathcal{L}$ . The algebras  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$  are defined as the quotient of  $\mathbf{k}B \ltimes \mathbf{k}\mathcal{L}$  by the two-sided ideal  $\mathfrak{J}$  generated by the elements  $\sigma^{m_H} - 1 - Q_s(\sigma)e_H$  where  $\sigma$  runs among the braided reflections of  $B$ ,  $s = \pi(\sigma)$  is the corresponding pseudo-reflection,  $H = \text{Ker}(s - 1)$ , and  $Q_s(X) = \sum_{k=0}^{m_H-1} a_{H,k} X^k - 1 \in \mathbf{k}[X]$  (see section 2.3.2 for more details).

Let  $\mathfrak{J}_H$  is the ideal of  $\mathbf{k}B$  generated by the  $\sigma^{m_H} - 1 - Q_s(\sigma) = \sigma^{m_H} - \sum_{k=0}^{m_H-1} a_{H,k} \sigma^k$ . This quotient  $H_{\mathbf{k}}(W) = (\mathbf{k}B)/\mathfrak{J}_H$  is by definition the Hecke algebra attached to  $W$  in the sense of Broué-Malle-Rouquier, and is the usual Iwahori-Hecke algebra of  $W$  when  $W$  is a finite Coxeter group. The following preliminary result explains the title, making our algebras appear as natural extensions of the Hecke algebra  $H_{\mathbf{k}}(W)$ .

**Proposition 1.1.** *Let  $\mathcal{L}$  be an admissible lattice for  $W$ . There exists a surjective algebra morphism  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L}) \rightarrow H_{\mathbf{k}}(W)$ , defining a split extension of  $H_{\mathbf{k}}(W)$ .*

*Proof.* The natural augmentation map  $\eta : \mathbf{k}\mathcal{L} \rightarrow \mathbf{k}$  defined by  $e_\lambda \mapsto 1$  induces surjective morphisms of  $\mathbf{k}$ -algebras  $\eta : \mathbf{k}B \ltimes \mathbf{k}\mathcal{L} \rightarrow \mathbf{k}B$  and  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L}) \rightarrow (\mathbf{k}B)/\mathfrak{J}_H = H_{\mathbf{k}}(W)$ . The splitting comes from the fact that the assumptions on  $\mathcal{L}$  imply that  $W$  belongs to  $\mathcal{L}$ , as the join of all the full cyclic subgroups. Then, the non-unital algebra morphism  $\mathbf{k}B \rightarrow \mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$  defined by  $b \mapsto be_W$  is easily checked to factorize through  $H_{\mathbf{k}}(W)$  and to provide a splitting.  $\square$

Our first result is a structure theorem of the following form, where the  $\mathbf{k}\tilde{H}_{x_*}$  are slight generalizations of the Hecke algebras attached to elements of  $\mathcal{L}$  and  $x_* \in \mathcal{L}$  is a representative of the orbit  $X \in \mathcal{L}/W$ .

**Theorem 1.2.** *There exists an isomorphism of  $\mathbf{k}$ -algebras*

$$\mathcal{C}_{\mathbf{k}}(W, \mathcal{L}) \simeq \bigoplus_{X \in \mathcal{L}/W} \text{Mat}_X(\mathbf{k}\tilde{H}_{x_*}).$$

When  $\mathcal{L}$  is the lattice of the reflection subgroups of a finite Coxeter group, the algebras  $\mathcal{C}(W, \mathcal{L})$  were introduced in [18], under the name  $\mathcal{C}_W$  and using a presentation by generators and relations, and proven to be generically semisimple. When  $W$  is the symmetric group,  $\mathcal{C}_W$  coincides with the diagram algebra of braids and ties of Aicardi and Juyumaya (see [1, 2]). Therefore the above theorem is a generalization of a theorem of Espinoza and Ryom-Hansen (see [14]), and was actually motivated by it. Note that, when  $W$  is the symmetric group, the lattice of parabolic subgroups coincides with the lattice of reflection subgroups.

We now return to the general case. We let  $K$  denote the field of fractions of  $R$  and  $\bar{K}$  an algebraic closure of  $K$ . The *BMR freeness conjecture* states that  $H_{\mathbf{k}}(W)$  is a free  $\mathbf{k}$ -module of rank  $|W|$ , and implies that  $H_{\mathbf{k}}(W)$  is generically semisimple. Up to extending the ring of definition  $R$  to a slightly larger Laurent polynomial ring  $R_u$ , an additional conjecture of Broué-Malle-Michel, which we recall in detail in section 3, states that  $H_{\mathbf{k}}(W)$  is a symmetric algebra when  $\mathbf{k}$  is a  $R_u$ -algebra, with a trace enjoying some uniqueness conditions. Of course both conjectures are true when  $W$  is a finite Coxeter group.

When  $\mathcal{L}$  is the lattice of parabolic subgroups of a finite complex reflection groups, the algebra  $\mathcal{C}(W, \mathcal{L})$  was introduced and called  $\mathcal{C}_W^p$  in [18]. It was conjectured there that  $\mathcal{C}_W^p$  is a free  $R$ -module

of rank  $|W| \times |\mathcal{L}_p|$ , where  $\mathcal{L}_p$  denotes the lattice of parabolic subgroups. A consequence of the above theorem is then the following one. We denote  $W_{x_*} < W$  the stabilizer of  $x_*$ .

**Theorem 1.3.** *The algebra  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$  is a free  $\mathbf{k}$ -module of finite rank if and only if the BMR freeness conjecture holds over  $\mathbf{k}$  for every  $x \in \mathcal{L}$  (this is in particular the case when  $W$  is a Coxeter group). In that case, its rank is  $|W| \times |\mathcal{L}|$ , and  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$  is semisimple when  $\mathbf{k}$  is an extension of  $K$ , and*

$$\mathcal{C}_{\bar{K}}(W, \mathcal{L}) \simeq \bar{K}W \ltimes \bar{K}\mathcal{L} \simeq \bigoplus_{X \in \mathcal{L}/W} \text{Mat}_X(\bar{K}W_{x_*}).$$

The BMR freeness conjecture is now proved for all irreducible reflection groups but the ones of Shephard-Todd types  $G_{17}$ ,  $G_{18}$  and  $G_{19}$  (see [3, 4, 17, 15, 20, 6, 7, 8]), therefore the above statement is actually almost unconditional, and reduces the proof of conjecture 5.10 in [18] to the original BMR freeness conjecture.

We finally (conditionnally) prove that these algebras are symmetric algebras. We call *strong freeness conjecture* for  $W$  the statement that  $H_R(W)$  admits a basis originating from elements of  $B$ . It turns out that the status of this conjecture is exactly the same as the original BMR freeness conjecture : for every group for which the BMR freeness conjecture has been proved so far, the proof provides a convenient basis.

**Theorem 1.4.** *Assume that the strong freeness conjecture as well as the Broué-Malle-Michel trace conjecture holds for all  $x \in \mathcal{L}$ . This is in particular the case if  $W$  is a finite Coxeter group. Then, for any commutative  $R_u$ -algebra  $\mathbf{k}$ , the algebra  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$  is a symmetric algebra.*

As an immediate corollary, we get that the diagram algebra of ‘braids and ties’ is a symmetric algebra as well.

**Acknowledgements.** I thank J.-Y. Hée and S. Bouc for discussions about root systems and lattices. I thank M. Calvez, A. Navas and J. Juyumaya for their invitation at the SUMA’16 conference in Valparaiso, where the original idea of this paper emerged.

## 2. STRUCTURE

**2.1. Semidirect extensions of group algebras by abelian algebras.** In this section, we first expose fairly general results, which are basically folklore, and which are needed in the sequel. To start with, the following proposition is an explicit version of what is known in the realm of the representation theory of finite groups as Mackey-Wigner’s method of “little groups” (see [23] §8.2). It can be seen as an explicit Morita equivalence (see [11] ex. 18.6). It is stated and proved in detail in [12], proposition 3.4, in the case  $G$  is finite. We explain below the additional arguments which are needed in the general case.

**Proposition 2.1.** *Let  $G$  be a group acting transitively (on the left) on a finite set  $X$ . Let  $\mathbf{k}$  be a commutative ring with 1, and let  $A$  be the  $\mathbf{k}$ -algebra  $G \ltimes \mathbf{k}^X$  where  $\mathbf{k}^X = \bigoplus_{x \in X} \mathbf{k}\epsilon_x$  is endowed with the product law  $(\epsilon_x \epsilon_{x'} = \delta_{x,x'} \epsilon_x)$  and the action of  $G$  is induced by the one on  $X$ . Then any choice of  $x_* \in X$  with stabilizer  $G_0 \subseteq G$  and any choice of a “section”  $\tau : X \rightarrow G$  such that  $\tau(x).x_* = x$  for all  $x \in X$ , define a unique isomorphism*

$$\theta : A \longrightarrow \text{Mat}_X(\mathbf{k}G_0)$$

sending each  $\epsilon_x \in \mathbf{k}^X$  ( $x \in X$ ) to  $\theta(\epsilon_x) := E_{x,x}$ , and each  $g \in G$  to

$$\theta(g) := \sum_{x \in X} \tau(gx)^{-1} g \cdot \tau(x) E_{gx,x}$$

(where  $E_{x,y} \in \text{Mat}_X(\mathbf{k})$  is the elementary matrix corresponding to  $x, y \in X$ ).

*Proof.* The proof given in proposition 3.4 of [12] that  $\theta$  is a surjective morphism does not use any finiteness assumption on  $G$ . It therefore remains to prove that  $\theta$  is injective. We prove this directly as follows. A  $\mathbf{k}$ -basis of  $A$  is given by the  $g\epsilon_\alpha$  for  $g \in G$  and  $\alpha \in X$ , and by definition

$$\theta(g\epsilon_\alpha) = \sum_{x \in X} \tau(gx)^{-1} g \tau(x) E_{gx,x} E_{\alpha,\alpha} = \tau(g\alpha)^{-1} g \tau(\alpha) E_{g\alpha,\alpha}.$$

It follows that a general linear combination  $\sum_{g,\alpha} \lambda_{g,\alpha} g \varepsilon_\alpha$  belongs to  $\text{Ker } \theta$  iff

$$0 = \sum_{g,\alpha} \lambda_{g,\alpha} \tau(g\alpha)^{-1} g \tau(\alpha) E_{g\alpha,\alpha}$$

which means that, for all  $\alpha \in X$ ,

$$\sum_{g \in G} \lambda_{g,\alpha} \tau(g\alpha)^{-1} g \tau(\alpha) E_{g\alpha,\alpha} = 0.$$

Let us fix such an  $\alpha \in X$ . For every  $\beta \in X$  we have

$$0 = \sum_{g \mid g.\alpha=\beta} \lambda_{g,\alpha} \tau(g\alpha)^{-1} g \tau(\alpha) E_{g\alpha,\alpha}$$

namely

$$0 = \tau(\beta)^{-1} \left( \sum_{g \mid g.\alpha=\beta} \lambda_{g,\alpha} g \right) \tau(\alpha)$$

which implies that, for all  $g \in G$ , we have  $\lambda_{g,\alpha} = 0$ . Since this holds for every  $\alpha \in X$  we get the conclusion.  $\square$

Let  $L$  be a join semilattice. That is, we have a finite partially ordered set  $L$  for which there exists a least upper bound  $x \vee y$  for every two  $x, y \in L$ . Let  $M$  be the semigroup with elements  $e_\lambda, \lambda \in L$  and product law  $e_\lambda e_\mu = e_{\lambda \vee \mu}$ . Such a semigroup is sometimes called a band.

If  $L$  is acted upon by some group  $G$  in an order-preserving way (that is  $x \leq y \Rightarrow g.x \leq g.y$  for all  $x, y \in L$  and  $g \in G$ ) then  $M$  is acted upon by  $G$ , so that we can form the algebra  $\mathbf{k}M \rtimes \mathbf{k}G$ . Up to exchanging meet and join, the algebra  $\mathbf{k}M$  is the Möbius algebra as in [24], definition 3.9.1. We recall from [18] a  $G$ -equivariant version of the classical isomorphism  $\mathbf{k}M \simeq \mathbf{k}^L$  of e.g. [24], theorem 3.9.2. Here  $\mathbf{k}^L$  is the algebra of  $\mathbf{k}$ -valued functions on  $L$ , that is the direct product of a collection indexed by the elements of  $L$  of copies of the  $\mathbf{k}$ -algebra  $\mathbf{k}$ . As before, to  $\lambda \in L$  we associate  $\varepsilon_\lambda \in \mathbf{k}^L$  defined by  $\varepsilon_\lambda(\lambda') = \delta_{\lambda,\lambda'}$  if  $\lambda' \in L$ .

**Proposition 2.2.** (see [18], proposition 3.9) *Let  $M$  be the band associated to a finite join semilattice  $L$ . For every commutative ring  $\mathbf{k}$ , the semigroup algebra  $\mathbf{k}M$  is isomorphic to  $\mathbf{k}^L$ . If  $L$  is acted upon by some group  $G$  as above, then  $\mathbf{k}M \rtimes \mathbf{k}G \simeq \mathbf{k}^L \rtimes \mathbf{k}G$ , the isomorphism being given by  $g \mapsto g$  for  $g \in G$  and  $e_\lambda \mapsto \sum_{\lambda \leq \mu} \varepsilon_\mu$ .*

By decomposing  $L$  as a disjoint union of  $G$ -orbits, by combining these two results one gets that  $\mathbf{k}M \rtimes \mathbf{k}G$  is isomorphic to a direct sum of  $|L/G|$  matrix algebras. This will turn out to be the main result from general algebra that is needed to prove our structure theorem.

**2.2. Braid groups of reflection subgroups.** Let  $W_0$  be a reflection subgroup of the reflection group  $W$ , and  $G$  a subgroup with  $W_0 < G < W$  normalizing  $W_0$ . For convenience we endow  $\mathbf{C}^n$  with a  $W$ -invariant unitary form.

The hyperplane complement associated to  $W$  is denoted  $X = \mathbf{C}^n \setminus \bigcup \mathcal{A}$ , and we let  $x_0 \in X$  denote the chosen base-point, so that  $B = \pi_1(X/W, x_0)$ . Let  $L \subset \mathbf{C}^n$  denote the fixed points set of  $W_0$ , namely the intersection of the set  $\mathcal{A}_L$  of all the reflecting hyperplanes associated to the reflections in  $W_0$ . Since  $G$  normalizes  $W_0$  we have  $g(L) = L$  for all  $g \in G$ . We let  $X_0 \subset L^\perp$  denote the hyperplane complement associated to  $W_0$  viewed as a reflection subgroup acting on  $L^\perp$ . We have  $X_0 = L^\perp \setminus \bigcup \mathcal{A}_L$ .

Let  $X^0 = \mathbf{C}^n \setminus \bigcup \mathcal{A}_L$ , and  $x_{00}$  the orthogonal projection of  $x_0$  on  $L^\perp$ . We write  $x_0 = x_1 + x_{00}$ , with  $x_1 \in L$ . Since  $x_0 \notin \bigcup \mathcal{A}_L$  we have  $x_{00} \in X_0$ , and the braid groups of  $W_0 < \text{GL}(L^\perp)$  can be defined as  $P_0 = \pi_1(X_0, x_{00})$ ,  $B_0 = \pi_1(X_0/W_0, x_{00})$ . The inclusion map  $(X_0, x_{00}) \subset (X^0, x_{00} + L)$  is a  $W_0$ -equivariant deformation retract through  $(z, t) \mapsto z_{L^\perp} + tz_L$  where  $z_{L^\perp}$  and  $z_L$  denote the orthogonal projections of  $z$  on  $L^\perp$  and  $L$ , respectively. Since  $x_{00} + L$  is retractable to  $x_0$ , it follows that this inclusion provides an isomorphism  $P_0 \simeq \pi_1(X^0, x_0)$  and, because of  $W_0$ -equivariance, an isomorphism  $B_0 \simeq \pi_1(X^0/W_0, x_0)$ .

Since  $W_0$  is normal inside  $G$ , the projection map  $X/W_0 \rightarrow X/G$  is a Galois covering, and we get a short exact sequence  $1 \rightarrow \pi_1(X/W_0, x_0) \rightarrow \pi_1(X/G, x_0) \rightarrow G/W_0 \rightarrow 1$ .

We consider the  $G$ -equivariant inclusion  $(X, x_0) \subset (X^0, x_0)$ . By standard arguments (see e.g. [13] proposition 2.2, or [5]) we know that the induced map  $P = \pi_1(X, x_0) \rightarrow \pi_1(X^0, x_0)$  is surjective, and that its kernel  $K$  is normally generated by the meridians around the hyperplanes in  $\mathcal{A}_L^c$ . Since the following diagram is commutative

$$\begin{array}{ccccccc}
 & & 1 & & 1 & & \\
 & & \downarrow & & \downarrow & & \\
 1 & \longrightarrow & K & \longrightarrow & \pi_1(X, x_0) & \longrightarrow & \pi_1(X^0, x_0) \longrightarrow 1 \\
 & & \parallel & & \downarrow & & \downarrow \\
 1 & \longrightarrow & K & \longrightarrow & \pi_1(X/W_0, x_0) & \longrightarrow & \pi_1(X^0/W_0, x_0) \longrightarrow 1 \\
 & & & & \downarrow & & \downarrow \\
 & & & & W_0 & \xlongequal{\quad} & W_0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 1 & & 1
 \end{array}$$

with the two columns and the top row being short exact sequences, it follows that the second row is exact and  $K = \text{Ker}(\pi_1(X/W_0, x_0) \rightarrow \pi_1(X^0/W_0, x_0))$ . Inside  $\pi_1(X/W_0, x_0)$ , the collection of meridians generating  $K$  become the collection of the elements  $\sigma^{m_\sigma}$  where  $\sigma$  runs among the collection of (distinguished) braided reflections around the hyperplanes in  $\mathcal{A}_L^c$  and  $m_\sigma$  is the order of their image in  $W_0 \subset W$ .

Since  $G$  stabilizes  $\mathcal{A}_L$ , the image of  $K$  under the injective map  $\pi_1(X/W_0, x_0) \rightarrow \pi_1(X/G, x_0)$  is a normal subgroup of  $\pi_1(X/G, x_0)$ , that we still denote  $K$ . We define the *generalized braid group associated to  $G$*  and denote  $B_G$  the quotient group  $\pi_1(X/G, x_0)/K$ .

Let us consider the projection map  $\pi : B \rightarrow W$ . By the above description,  $B_G$  is the quotient of  $\hat{B}_G = \pi_1(X/G, x_0) = \pi^{-1}(G)$  by  $K$ , and the short exact sequence  $1 \rightarrow \pi_1(X/W_0) \rightarrow \pi_1(X/G, x_0)/K \rightarrow G/W_0 \rightarrow 1$  induces a short exact sequence  $1 \rightarrow \pi_1(X/W_0)/K \rightarrow B_G \rightarrow G/W_0 \rightarrow 1$ . Identifying  $\pi_1(X/W_0)/K$  with  $\pi_1(X^0/W_0, K)$  we get a short exact sequence  $1 \rightarrow B_0 \rightarrow B_G \rightarrow G/W_0 \rightarrow 1$ .

We now consider the central element  $\pi_0 \in P_0$  defined as the class inside  $P_0 = \pi_1(X_0, x_{00})$  of the loop  $\gamma_0(t) = x_{00} \exp(2i\pi t)$ . By the above identifications, it is identified inside  $\pi_1(X^0, x_0)$  with the path  $\gamma_1 \star \gamma_0 \star \gamma_1^{-1}$ , where  $\gamma_1(t) = x_{00} + tx_1$  (recall that  $x_0 = x_1 + x_{00}$ ). We prove that it remains a central element inside  $B_G = \pi_1(X/G, x_0)/K$ .

For this, let us consider a path  $\gamma : x_0 \rightsquigarrow g.x_0$  inside  $X$ . We need to prove that the composite  $\gamma^{-1} \star (g.\gamma_1 \star g.\gamma_0 \star g.\gamma_1^{-1})^{-1} \star \gamma \star (\gamma_1 \star \gamma_0 \star \gamma_1^{-1})$ , which is a path  $x_0 \rightsquigarrow x_0$  inside  $X$ , belongs to  $K$ . This means that its class must be 0 inside  $\pi_1(X, x_0)/K = \pi_1(X^0, x_0)$ . Therefore we need to prove that  $\gamma \star \gamma_1 \star \gamma_0 \star \gamma_1^{-1} : x_0 \rightsquigarrow g.x_0$  is homotopic to  $g.\gamma_1 \star g.\gamma_0 \star g.\gamma_1^{-1} \star \gamma$  inside  $X^0$ . For this, consider the following map  $\tilde{H} : [0, 3] \times [0, 1] \rightarrow X^0$  defined, for  $t, u \in [0, 1]$ , by  $\tilde{H}(t, u) = \gamma(u)_{L^\perp} + (1-t)\gamma(u)_L$ ,  $\tilde{H}(1+t, u) = \gamma(u)_{L^\perp} \exp(2i\pi t)$ ,  $\tilde{H}(2+t, u) = \gamma(u)_{L^\perp} + t\gamma(u)_L$ . It is not difficult to check that indeed  $\tilde{H}(t, u) \in X^0$  for all  $t, u$ , and that  $\tilde{H}$  is continuous. Moreover, the boundary of the rectangle  $[0, 3] \times [0, 1]$  has for image the union of the two paths we are interested in. It follows that these two paths are homotopic, which proves our claim.

### 2.3. Proof of the structure theorem.

**2.3.1. Generalized Hecke algebras.** We now attach to an admissible lattice  $\mathcal{L}$  the following datas. To each  $x \in \mathcal{L}$  we attach

- the ring  $R_x = \mathbf{Z}[a_{H,i}, a_{H,0}^{\pm 1}]$  where  $H$  runs among all  $H \in x$ , and  $1 \leq i \leq m_H - 1$

- the stabilizer  $G_x < W$  of  $x \in \mathcal{L}$  and the group  $\hat{B}_x = \hat{B}_{G_x} = \pi^{-1}(G_x)$  associated to  $W_0 < G_x < N_W(W_0)$ , where  $W_0$  is the full reflection subgroup associated to  $x \in \mathcal{L}$ .
- the group  $B_x = B_{G_x}$  as in the previous section.

The generalized Hecke algebra  $\tilde{H}_x$  associated to  $x \in \mathcal{L}$  is then defined as the quotient of the group algebra  $R_x B_x$  by the ideal generated by the Hecke relations  $\sigma^{m_H} - \sum a_{H,i} \sigma^i$  for  $\sigma$  a braided reflection with respect to an hyperplane in  $x$ . Equivalently, it is the quotient of the group algebra  $R_x \hat{B}_x$  by the relations  $\sigma^{m_H} - \sum a_{H,i} \sigma^i$  for  $\sigma$  a braided reflection with respect to an hyperplane of  $x$  and  $\sigma^{m_H} = 1$  for  $\sigma$  a braided reflection with respect to an hyperplane of  $\mathcal{A} \setminus x$ .

Now recall the short exact sequence  $1 \rightarrow B_0 \rightarrow B_x \rightarrow G/W_0 \rightarrow 1$ , and consider the induced injective map  $R_x B_0 \rightarrow R_x B_x$ . We let  $\mathfrak{h}_0$  denote the ideal of  $R_x B_0$  generated by the  $\sigma^{m_H} - \sum a_{H,i} \sigma^i$  for  $\sigma$  a braided reflection with respect to an hyperplane of  $x$ . By definition the quotient algebra  $H_0 = R_x B_0 / \mathfrak{h}_0$  the usual Hecke algebra associated to  $W_0$ . We let  $\mathfrak{h}_x$  the ideal of  $R_x B_x$  generated by the same elements, and choose a system  $b_1, \dots, b_m$  of representatives inside  $B_x$  of  $B_x / B_0 \simeq G/W_0$ . Since the generating set of  $\mathfrak{h}_x$  is stable under  $B_x$ -conjugation, we have  $\mathfrak{h}_x = \bigoplus_{i=1}^m b_i \mathfrak{h}_0$ . This implies that, as a right  $R_x B_0$ -module,  $\tilde{H}_x = \bigoplus_{i=1}^m (b_i (R_x B_0)) / (b_i \mathfrak{h}_0) = \bigoplus_{i=1}^m (b_i (R_x B_0)) / (b_i \mathfrak{h}_0)$ . Now,  $(b_i (R_x B_0)) / (b_i \mathfrak{h}_0)$  contains (the class of)  $b_i$  and is clearly a free  $H_0$ -module of rank 1. This proves that  $\tilde{H}_x = R_x B_x / \mathfrak{h}_x$  is a free  $H_0$ -module of rank  $|G/W_0|$ . In particular,  $\tilde{H}_x$  is a free  $R_x$ -module of rank  $|G|$  if and only if  $H_0$  is a free  $R_x$ -module of rank  $|W_0|$ . This latter assumption is exactly the BMR freeness conjecture for  $W_0$ .

**2.3.2. Image of the defining ideal.** Let  $\mathcal{L}$  be an admissible lattice. The group  $B$  acts on  $\mathcal{L}$  via the natural projection map  $B \rightarrow W$ . We denote  $\mathfrak{J}$  the ideal of  $\mathbf{k}B \ltimes \mathbf{k}\mathcal{L}$  generated by the elements  $\sigma^m - 1 - Q_s(\sigma)e_H$  where

- $s$  runs among the distinguished pseudo-reflections of  $W$ ,
- $\sigma$  is a braided reflection attached to it,
- $H = \text{Ker}(s - 1)$  is the fixed hyperplane, and
- $e_H \in \mathbf{k}\mathcal{L}$  is the idempotent attached to  $\{H\} \in \mathcal{L}$
- $m$  is the order of  $s$
- $Q_s(X) = \sum_{k=0}^{m-1} a_{H,k} X^k - 1$ , where  $\prod_{k=1}^m (X - u_{s,i}) = X^m + \sum_{k=0}^{m-1} a_{H,k} X^k$ .

Let  $s$  be a reflection, and  $m$  its order. Let  $1 \leq k < m$ . For any hyperplane  $H \in \mathcal{A}$ , we have  $s(H) = H \Leftrightarrow s^k(H) = H$ . It follows that, for every  $x \in \mathcal{L}$ , we have  $s.x = x \Leftrightarrow s^k.x = x$ . We consider the composite  $\theta$  of the maps provided by propositions 2.2 and 2.1

$$\mathbf{k}B \ltimes \mathbf{k}\mathcal{L} \rightarrow \mathbf{k}B \ltimes \mathbf{k}\mathcal{L} \rightarrow \mathbf{k}B \ltimes \mathbf{k}^X \rightarrow \text{Mat}_X(\mathbf{k}\hat{B}_{x_*})$$

where  $\hat{B}_{x_*} = \hat{B}_{G_{x_*}} = \pi^{-1}(G_{x_*})$  is the stabilizer of  $x_* \in \mathcal{L}$  and  $X$  is the orbit of  $x_*$  under  $B$  (or  $W$ ). We have, for all  $r \in \mathbf{Z}$ ,

$$\theta(e_H) = \sum_{\substack{x \in X \\ H \in x}} E_{x,x} \quad \text{and} \quad \theta(\sigma^r) = \sum_{x \in X} \tau(s^r.x)^{-1} \sigma^r \tau(x) E_{s^r.x, x}.$$

Since  $H \in x \Rightarrow s^r.x = x$ , this implies

$$\theta(\sigma^r e_H) = \sum_{\substack{x \in X \\ H \in x}} \tau(x)^{-1} \sigma^r \tau(x) E_{x,x} \quad \text{and} \quad \theta(Q_s(\sigma)e_H) = \sum_{\substack{x \in X \\ H \in x}} \tau(x)^{-1} Q_s(\sigma) \tau(x) E_{x,x}$$

hence the image under  $\theta$  of  $\sigma^m - 1 - Q_s(\sigma)e_H$  is equal to

$$\sum_{\substack{x \in X \\ H \notin x}} \tau(x)^{-1} (\sigma^m - 1) \tau(x) E_{x,x} + \sum_{\substack{x \in X \\ H \in x}} \tau(x)^{-1} (\sigma^m - 1 - Q_s(\sigma)) \tau(x) E_{x,x}.$$

Now recall the elementary fact that, for any ring  $A$  with 1 (commutative or not), the twosided ideal of the matrix algebra  $\text{Mat}_N(A)$  generated by a collection  $S^\alpha, \alpha \in F$  of matrices  $S^\alpha = (S_{i,j}^\alpha)_{1 \leq i,j \leq N}$  is equal to  $\text{Mat}_N(I)$  where  $I$  is the twosided ideal of  $A$  generated by the  $S_{i,j}^\alpha$  for  $\alpha \in F, 1 \leq i,j \leq N$ . It follows that image of the ideal  $\mathfrak{J}$  inside  $\text{Mat}_X(\mathbf{k}\hat{B}_{x_*})$  is  $\text{Mat}_X(\mathfrak{J}_X)$  where  $\mathfrak{J}_X$  is the ideal of  $\mathbf{k}\hat{B}_{x_*}$  generated by the  $\sigma_x^m - 1$  for  $H \notin x, x \in X$  and the  $\sigma_x^m - 1 - Q_s(\sigma_x)$

for  $H \in x, x \in X$ , where  $\sigma_x = \tau(x)^{-1}\sigma\tau(x)$ . This is the same as the ideal of  $\mathbf{k}\hat{B}_{x_*}$  generated by the  $\sigma^m - 1$  for  $\sigma$  a braided reflection around some  $H \notin x_*$ , and the  $\sigma^m - 1 - Q_s(\sigma)$  and  $\sigma$  for a braided reflection around some  $H \in x_*$ . Therefore  $\mathbf{k}\hat{B}_{x_*}/\mathfrak{J}_X = \mathbf{k}\tilde{H}_x$  whence, from the isomorphism  $\mathbf{k}B \rtimes \mathbf{k}\mathcal{L} \simeq \bigoplus_{X \in \mathcal{L}/W} \text{Mat}_X(\mathbf{k}\hat{B}_{x_*})$  we get the following.

**Theorem 2.3.** *Let  $\mathcal{L}$  be an admissible lattice. Then we have an isomorphism*

$$\mathcal{C}_{\mathbf{k}}(W, \mathcal{L}) \simeq \bigoplus_{X \in \mathcal{L}/W} \text{Mat}_X(\mathbf{k}\tilde{H}_{x_*}).$$

The following corollary completes the proof of theorem 1.3.

**Corollary 2.4.** *The algebra  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$  is a free  $\mathbf{k}$ -module of finite rank if and only if the BMR freeness conjecture holds over  $\mathbf{k}$  for every  $x \in \mathcal{L}$ . In that case, its rank is  $|W| \times |\mathcal{L}|$ , and it is generically semisimple.*

The fact that it is generically semisimple is a consequence of the fact that, under the specialization morphism  $\varphi : R \rightarrow \mathbf{Q}$  defined by  $a_{H,i} \mapsto 0$  if  $i > 0$ ,  $a_{H,0} \mapsto 0$ , the algebra  $\mathcal{C}(W, \mathcal{L}) \otimes_{\varphi} \mathbf{Q}$  becomes isomorphic to a semidirect product  $\mathbf{Q}W \rtimes \mathbf{Q}\mathcal{L} \simeq \bigoplus_{\mathcal{L}/W} \text{Mat}_X(\mathbf{Q}W_{x_*})$ , where  $W_{x_*} < W$  is the stabilizer of  $x_* \in \mathcal{L}$ . By Maschke's theorem we get that  $\mathcal{C}(W, \mathcal{L}) \otimes_{\varphi} \mathbf{Q}$  is semisimple, and therefore  $\mathcal{C}(W, \mathcal{L})$  is generically semisimple as soon as it is a free  $R$ -module of finite rank. By Tits' deformation theorem we get that

$$\mathcal{C}_{\bar{K}}(W, \mathcal{L}) \simeq \bar{K}W \rtimes \bar{K}\mathcal{L} \simeq \bigoplus_{\mathcal{L}/W} \text{Mat}_X(\bar{K}W_{x_*}).$$

Since the BMR freeness conjecture is now proved for all irreducible reflection groups but the ones of Shephard-Todd types  $G_{17}$ ,  $G_{18}$  and  $G_{19}$  (see [3, 4, 17, 15, 20, 6, 7, 8]), this proves the following.

**Corollary 2.5.** *The algebra  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$  is a free  $\mathbf{k}$ -module of rank  $|W| \times |\mathcal{L}|$ , and is generically semisimple, except possibly if there exists  $x \in \mathcal{L}$  whose associated reflection group admits an irreducible component of Shephard-Todd type  $G_{17}$ ,  $G_{18}$  or  $G_{19}$ .*

### 3. TRACES

In this section, we slightly extend the ring of definition, for convenience. For  $W$  a given complex reflection group, we denote  $R_u = \mathbf{Z}[u_{c,i}^{\pm 1}]$ , where  $c$  runs among the conjugacy classes of distinguished pseudo-reflections, and  $i$  between 1 and the order of (a representative of)  $c$ . We consider  $R$  as a subring of  $R_u$  where  $a_{H,i}, H \in \mathcal{A}$  is mapped to the  $(m_H - i)$ -th symmetric function in the  $u_{c,k}$ , where  $c$  is the conjugacy class corresponding to the distinguished pseudo-reflection with hyperplane  $H$ . We let  $H_u$  denote the Hecke algebra of  $W$  defined over  $R_u$ , that is  $H_u = H(W) \otimes_R R_u$ .

**3.1. Reminder on canonical traces.** Let  $W$  be a complex reflection group,  $B$  its braid group,  $H = H_u$  its Hecke algebra, defined over the ring of definition  $R_u = \mathbf{Z}[u_{c,i}^{\pm 1}]$ . Let  $x \mapsto \bar{x}$  the automorphism of  $R_u$  defined by  $u_{c,i} \mapsto u_{c,i}^{-1}$ . The group antiautomorphism  $g \mapsto g^{-1}$  on  $B$  induces an antiautomorphism of  $\mathbf{Z}$ -algebras  $\mathbf{a} : R_u B \rightarrow R_u B$  such as  $\mathbf{a}(\lambda g) = \bar{\lambda} g^{-1}$  for all  $\lambda \in R_u$  and  $g \in B$ . The Hecke ideal  $\mathfrak{J}_H$  of  $R_u B$  is stable by  $\mathbf{a}$  hence  $\mathbf{a}$  induces an automorphism of  $H$ . It has the property that, for all parabolic subalgebras  $H_0$  of  $H$ ,  $H_0$  is  $\mathbf{a}$ -stable and the restriction of  $\mathbf{a}$  to  $H_0$  coincides with the antiautomorphism associated to  $H_0$ . Let  $t : H \rightarrow R$  be a linear form. We assume that  $H$  admits a  $R_u$ -basis whose elements are (images of) elements of  $B$ . Note that this is proved so far for all complex reflection groups but the ones having an irreducible component of type  $G_{17}$ ,  $G_{18}$  or  $G_{19}$ . We denote  $\pi$  the natural central element of  $P = \text{Ker}(B \rightarrow W)$ . We consider the following assumptions on  $t$ .

- (1)  $t$  is a symetrizing trace on  $H$ .
- (2) The trace induced on the specialization  $\mathbf{C}W$  of  $H$  is the usual trace on the group algebra  $\mathbf{C}W$
- (3) For all  $h \in H$ , we have  $\overline{t(\mathbf{a}(h))}t(\pi) = t(h\pi)$ .

In [9] proposition 2.2 it is proven that, if there exists a trace satisfying these assumptions, then it is unique. It is also proven there that, in case  $W$  is a Coxeter group, then the trace given by  $t(T_w) = 0$  if  $w \neq 1$ ,  $t(T_1) = 1$ , satisfies these assumptions.

**3.2. Traces on generalized Hecke algebras.** Let  $\mathcal{L}$  be an admissible lattice, and  $x \in \mathcal{L}$ . Let  $W_0$  denote the full reflection subgroup attached to  $x$  and  $H_0$  the corresponding Hecke algebra. We already proved that the generalized Hecke algebra  $\tilde{H}_x$  attached to  $x$  is a free  $H_0$ -module of the form  $\bigoplus_{i=1}^m b_i H_0$  where the  $b_i$  are (classes inside  $\tilde{H}_x$  of) representatives of  $B_x/B_0 \simeq G_x/W_0$ . Obviously one can assume  $b_1 = 1$  hence  $b_1 H_0 = H_0$ . Assume that we are given a trace  $t : H_0 \rightarrow R_u$  satisfying the conditions of the previous section. We extend it as a linear form  $t : \tilde{H}_x \rightarrow R_u$  by  $t(b_i H_0) = 0$  if  $i > 1$ .

**Proposition 3.1.** *The extended linear form  $t : \tilde{H}_x \rightarrow R_u$  is a symmetrizing trace.*

*Proof.* In order for it to be a trace one needs to check that for all  $a_1, a_2 \in H_0$  and  $i, j$  we have  $t(b_i a_1 b_j a_2) = t(b_j a_2 b_i a_1)$ . But clearly both terms are 0 if  $b_j \notin b_i^{-1} H_0$ . Therefore we need to check that  $t(b_i a_1 b_i^{-1} a_2) = t(b_i^{-1} a_2 b_i a_1)$  for all  $i$  and  $a_1, a_2 \in H_0$ . But this means  $t(b_i a_1 (b_i^{-1} a_2 b_i) b_i^{-1}) = t(b_i^{-1} a_2 b_i a_1)$ . Since  $a_2 \mapsto b_i^{-1} a_2 b_i$  induces a bijection of  $H_0 \hookrightarrow \tilde{H}_x$  this is equivalent to saying that  $t(b_i a_1 a_2 b_i^{-1}) = t(a_2 a_1)$  for all  $a_1, a_2 \in H_0$ . But  $t(a_2 a_1) = t(a_1 a_2)$  whence we need to check that, for all  $i$  and all  $a \in H_0$ , we have  $t(b_i a b_i^{-1}) = t(a)$ . This holds true for the following reason. Let  $b \in B_x$ , and consider the map  $a \mapsto t(b a b^{-1})$ . This is a trace on  $H_0$ , which satisfies obviously the conditions (1) and (2) of the previous section. It also satisfies condition (3) if we can prove that  $b \pi_0 b^{-1} = \pi_0$  where  $\pi_0$  is the natural central element of the pure braid group  $P_0$  of  $W_0$ . But this was proven in section 2.2 above. Therefore  $t$  is a trace on  $\tilde{H}_x$ . Taking a basis  $e_1, \dots, e_N$  of  $H_0$  and letting  $e'_1, \dots, e'_N$  its dual basis, so that  $t(e_i e'_j) = \delta_{ij}$ , we get that the  $b_i e_j$  form a basis for  $\tilde{H}_x$ , with dual basis  $e'_j b_i^{-1}$ . Indeed,  $t(b_i e_j e'_r b_s^{-1}) = t(b_s^{-1} b_i e_j e'_r) = 0$  unless  $i \neq s$ , and in that case it is equal to  $t(e_j e'_r) = \delta_{jr}$ . Therefore  $t$  is a symmetrizing trace.  $\square$

**3.3. Symmetrizing trace.** We recall the following standard property of traces on matrix algebras, the proof being easy and left to the reader.

**Lemma 3.2.** *Let  $\mathbf{k}$  be a commutative ring with 1,  $A$  a  $\mathbf{k}$ -algebra and  $N \geq 1$ . There is a 1-1 correspondence between trace forms on  $A$  and trace forms on  $\text{Mat}_N(A) = \text{Mat}_N(\mathbf{k}) \otimes_{\mathbf{k}} A$ , the correspondence being given by  $t \mapsto \text{tr} \otimes t$ , where  $\text{tr} : \text{Mat}_N(\mathbf{k}) \rightarrow \mathbf{k}$  is the matrix trace. Moreover  $\text{tr} \otimes t : \text{Mat}_N(\mathbf{k}) \otimes_{\mathbf{k}} A \rightarrow \mathbf{k} \otimes_{\mathbf{k}} \mathbf{k} = \mathbf{k}$  is symmetrizing if and only if  $t$  is symmetrizing.*

From the isomorphism  $(\mathbf{k}B \ltimes \mathbf{k}\mathcal{L})/\mathfrak{J} \simeq \bigoplus_{X \in \mathcal{L}/W} \text{Mat}_X(\mathbf{k}\tilde{H}_{x_*})$  we are able to construct a trace form, as

$$\bigoplus_{X \in \mathcal{L}/W} t_{x_*} \otimes \text{tr} : \bigoplus_{X \in \mathcal{L}/W} \text{Mat}_X(\mathbf{k}\tilde{H}_{x_*}) = \bigoplus_{X \in \mathcal{L}/W} \text{Mat}_X(\mathbf{k}) \otimes_{\mathbf{k}} \mathbf{k}\tilde{H}_{x_*} \rightarrow \mathbf{k}$$

and by the above property it is a symmetrizing form. This proves the following.

**Theorem 3.3.** *Let  $\mathcal{L}$  be an admissible lattice for  $W$ , and  $\mathbf{k}$  a commutative  $R_u$ -algebra. If the Broué-Malle-Michel trace conjecture holds for all  $x \in \mathcal{L}$ , then the algebra  $\mathcal{C}_{\mathbf{k}}(W, \mathcal{L})$  is a symmetric algebra. It is in particular the case when  $W$  is a real reflection group.*

#### 4. MAIN EXAMPLES

We recall that a reflection subgroup  $W_0$  of  $W$  is called *full* if, for every reflection  $s \in W_0$ , all the reflections with respect to the same reflecting hyperplane belong to  $W_0$ . Such a reflection subgroup is uniquely determined by the set of its reflecting hyperplanes. Of course reflection subgroups of real reflection groups and, more generally, of 2-reflection groups, are full.

Let  $\mathcal{L}_{\infty}$  denote the poset of all full reflection subgroups, ordered by inclusion. For convenience, we prefer to consider it as a poset of subsets of  $\mathcal{L}$ , also ordered by inclusion.

Recall that a subset  $\mathcal{L} \subset \mathcal{L}_{\infty}$  is called *admissible* if it is a sub-join-semilattice of  $\mathcal{L}_{\infty}$  which satisfies the following conditions:



- (1) It is  $W$ -stable
- (2) It contains all  $\{H\}$ , for  $H \in \mathcal{A}$ , as well as the trivial subgroup.

Because such an  $\mathcal{L}$  always contains a minimal element (the trivial group), there is no ambiguity in the definition of the semi-lattice : the fact that  $a \vee b$  exists for every two elements of  $\mathcal{L}$  is in this case equivalent to saying that every finite subset of elements, including the empty one, admits a join. Moreover, since such an  $\mathcal{L}$  is always finite, it is automatically a lattice. Therefore, we can equivalently talk about admissible *lattices*.

**4.1. The category of admissible semi-lattices and maps.** Let  $\mathcal{L}$  and  $\mathcal{L}'$  be two admissible semi-lattices. A map  $\mathcal{L} \rightarrow \mathcal{L}'$  is called admissible if it is a  $W$ -equivariant morphism of join semi-lattices which is the identity on the cyclic and trivial reflection subgroups. The collection of admissible semi-lattices with morphisms the admissible maps forms a (small, finite) category  $\mathcal{CL}_W$ , and  $\mathcal{C}_k(W, \bullet)$  defines a functor from  $\mathcal{CL}_W$  to the category of (associative, unital)  $\mathbf{k}$ -algebras. The category  $\mathcal{CL}_W$  admits a terminal object that we call  $\mathcal{L}_2$  : it is the subset of  $\mathcal{L}_\infty$  made of the trivial and cyclic reflection subgroups together with the whole group  $W$ . Obviously, for every admissible  $\mathcal{L}$  there exists exactly one admissible map  $\mathcal{L} \rightarrow \mathcal{L}_2$ . In particular there exists exactly one admissible map  $\mathcal{L}_\infty \rightarrow \mathcal{L}_2$ .

More generally, define the parabolic rank of a reflection subgroup  $W_0$  as the rank of the smallest parabolic subgroup containing  $W_0$ , or equivalently as the codimension of its set of fixed points. Then, the sub-poset  $\mathcal{L}_n$  made of all reflection subgroups of parabolic rank at most  $n$  plus the whole group is an admissible semi-lattice as soon as  $n \geq 2$ , and there is an admissible map  $\mathcal{L}_m \rightarrow \mathcal{L}_n$  when  $m \geq n$  given by  $W_0 \mapsto W_0$  if  $W_0$  has parabolic rank at most  $n$ , and  $W_0 \mapsto W$  if  $W_0$  has rank at least  $n + 1$ . This applies to  $m = \infty$  as well.

**4.2. The semi-lattice  $\mathcal{L}_2$ .** The  $W$ -orbits of  $\mathcal{L}_2$  are  $\{\{1\}\}$ ,  $\{W\}$  together with the  $b_c = \{\{H\}; H \in c\}$  for every  $c \in \mathcal{A}/W$ . It is immediately checked that  $\mathbf{k}\tilde{H}_1 = \mathbf{k}W$  and  $\mathbf{k}\tilde{H}_{W_*} = H_{\mathbf{k}}(W)$ . From theorem 2.3 we get that

$$\mathcal{C}_k(W, \mathcal{L}_2) = \mathbf{k}W \oplus H_{\mathbf{k}}(W) \oplus \bigoplus_{c \in \mathcal{A}/W} \text{Mat}_{|c|}(\mathbf{k}\tilde{H}_{c_*})$$

A remarkable fact about the  $x = \{H\} \in \mathcal{L}$  of rank 1, for any admissible poset, is that the generalized Hecke algebras  $\tilde{H}_x$  are free deformations of the group algebra  $\mathbf{k}G(H)$ , where  $G(H) = \{w \in W \mid w(H) = H\}$ , without having to invoke the BMR freeness conjecture (or, said differently, it corresponds to the trivial case (rank 1) of the BMR freeness conjecture).

**4.3. The case of finite Coxeter groups.** Assume that  $W$  is a real reflection group, and let  $(W, S)$  be a Coxeter system attached to it. Then  $B$  admits a presentation as an Artin group, with generators  $b_s, s \in S$ . The map  $B \rightarrow W$  admits a natural set-theoretic section, called Tits' section, and defined by  $w \mapsto b_w = b_{s_1} \dots b_{s_n}$  where  $s_i \in S$  and  $w = s_1 \dots s_n$  is an expression of  $w$  as a product of the generators of minimal length. The classical theory tells us that it is well-defined. We denote  $g_w$  the image of  $b_w$  inside  $\mathcal{C}(W, \mathcal{L})$  under the natural  $R$ -algebra morphism  $RB \rightarrow \mathcal{C}(W, \mathcal{L})$ .

Since the BMR freeness conjecture is true for all reflection subgroups of  $W$ , from theorem 1.3 we know that  $\mathcal{C}(W, \mathcal{L})$  is a free  $R$ -module of rank  $|W| \times |\mathcal{L}|$ . More precisely, we have the following.

**Proposition 4.1.** *Let  $W$  be a finite Coxeter group and  $\mathcal{L}$  an admissible lattice. Then  $\mathcal{C}(W, \mathcal{L})$  admits for basis the elements  $g_w e_L$  for  $w \in W$  and  $L \in \mathcal{L}$ .*

*Proof.* Since the collection  $\{g_w e_L; w \in W, L \in \mathcal{L}\}$  has the right cardinality, it is sufficient to prove that it spans the free  $R$ -module of finite rank  $\mathcal{C}(W, \mathcal{L})$ . For this we consider its span that we denote  $V$  ; we remark that  $1 \in V$ , and prove that it is a left ideal of the  $R$ -algebra  $\mathcal{C}(W, \mathcal{L})$ . Since the  $g_s, s \in S$  and  $e_L, L \in \mathcal{L}$  generate  $RB \rtimes R\mathcal{L}$  as an algebra, they also generate  $\mathcal{C}(W, \mathcal{L})$  and therefore it is sufficient to show that  $g_s.x \in V$  and  $e_L.x \in V$  for  $x$  running among a spanning set of  $V$ . Setting  $x = g_w e_M$  for some  $w \in W, M \in \mathcal{L}$ , we get  $e_L g_w e_M = g_w e_{w^{-1}(L)} e_M = g_w e_{w^{-1}(L) \vee M} \in V$ . Let  $\ell : W \rightarrow \mathbf{N} = \mathbf{Z}_{\geq 0}$  denote the classical length function. If  $\ell(sw) = \ell(w) + 1$ , then  $g_s x = g_s g_w e_M = g_{sw} e_M \in V$ . If not,  $w$  can be written  $w = sw'$  with  $\ell(w') = \ell(w) - 1$ . Then

$g_s g_w = g_s^2 g_{w'} = g_{w'} + (u_s - 1)e_{\langle s \rangle}(1 + g_s)g_{w'} = g_{w'} + (u_s - 1)e_{\langle s \rangle}g_{w'} + (u_s - 1)e_{\langle s \rangle}g_s g_{w'} = g_{w'} + (u_s - 1)g_{w'}e_{\langle s w' \rangle} + (u_s - 1)e_{\langle s \rangle}g_w$ , hence  $g_s g_w e_M = g_{w'}e_M + (u_s - 1)g_{w'}e_{\langle s w' \rangle}e_M + (u_s - 1)e_{\langle s \rangle}g_w e_M = g_{w'}e_M + (u_s - 1)g_{w'}e_{\langle s w' \rangle \vee M} + (u_s - 1)e_{\langle s \rangle}g_w e_M \in V$ . This proves the claim.  $\square$

This proposition implies the following corollary, which could also be directly obtained from the approach of [18] – for instance by extending the left action of  $\mathcal{C}_W$  on itself to an action of  $\mathcal{C}(W, \mathcal{L}_\infty)$ .

**Corollary 4.2.** *If  $W$  is a finite Coxeter group, then  $\mathcal{C}_W \simeq \mathcal{C}(W, \mathcal{L}_\infty)$ .*

*Proof.* The elements  $g_s, s \in S$  and  $e_H, H \in \mathcal{A}$  clearly satisfy inside  $\mathcal{C}(W, \mathcal{L}_\infty)$  the defining relations of  $\mathcal{C}_W$ , and from this we get an algebra morphism  $\mathcal{C}_W \rightarrow \mathcal{C}(W, \mathcal{L}_\infty)$ . From the above proposition and theorem 3.4 of [18] we get that it maps a basis of  $\mathcal{C}_W$  to a basis of  $\mathcal{C}(W, \mathcal{L}_\infty)$ , and therefore it is an isomorphism.  $\square$

Therefore, the construction of  $\mathcal{C}(W, \mathcal{L}_\infty)$  indeed generalizes to the complex reflection group case the algebra  $\mathcal{C}_W$  of a finite Coxeter group introduced in [18].

**4.4. The parabolic lattice.** A  $W$ -stable subposet of  $\mathcal{L}_\infty$  is given by the collection  $\mathcal{L}_p$  of parabolic subgroups. It can be identified with the arrangement lattice  $L(\mathcal{A})$ , that is the collection of all intersections of hyperplanes in  $\mathcal{A}$ , ordered by reverse inclusion. More precisely, there exists a map  $\text{Fix} : \mathcal{L} \rightarrow L(\mathcal{A})$  where  $\text{Fix}(x)$  is the intersection of all reflecting hyperplanes in  $x$ , and its restriction to  $\mathcal{L}_p$  is a bijection.

**Proposition 4.3.** *For  $x \in \mathcal{L}_\infty$  a reflection subgroup, let  $[x] \in \mathcal{L}_p$  denote the parabolic closure of  $x$ . Then  $x \mapsto [x]$  is an admissible map  $\mathcal{L}_\infty \rightarrow \mathcal{L}_p$  inducing a quotient map  $\mathcal{C}(W, \mathcal{L}_\infty) \rightarrow \mathcal{C}(W, \mathcal{L}_p)$ .*

*Proof.* First note that, for every  $E, F \subset W$ , we have  $\text{Fix}(E \cup F) = \text{Fix}(E) \cap \text{Fix}(F)$ ,  $\text{Fix}(E) = \text{Fix}(\langle E \rangle)$ , and  $\text{Fix}(x) = \text{Fix}([x])$  if  $x$  is a reflection subgroup. From this we get that, for all  $x, y \in \mathcal{L}$ , we have on the one hand  $\text{Fix}(\langle [x], y \rangle) = \text{Fix}(\langle x \cup y \rangle) = \text{Fix}(x \cup y) = \text{Fix}(x) \cap \text{Fix}(y)$ , and on the other hand  $\text{Fix}(\langle [x] \cup [y] \rangle) = \text{Fix}(\langle [x] \cup [y] \rangle) = \text{Fix}([x] \cup [y]) = \text{Fix}([x]) \cap \text{Fix}([y]) = \text{Fix}(x) \cap \text{Fix}(y)$ . Since  $\text{Fix}$  is a bijection  $\mathcal{L}_p \rightarrow L(\mathcal{A})$  this proves  $\langle [x], y \rangle = \langle [x] \cup [y] \rangle$ , and this proves the claim, the  $W$ -invariance being obvious.  $\square$

From this we recover the definition of  $\mathcal{C}_W^p = \mathcal{C}(W, \mathcal{L}_p)$  given in [18] in the case of a finite Coxeter group, and extend the map  $\mathcal{C}(W, \mathcal{L}_\infty) \rightarrow \mathcal{C}_W^p$  to the complex reflection group case.

**4.5. Root systems.** Let  $R$  be a reduced root system (in the sense of Bourbaki),  $W$  the associated real reflection group. To each  $\alpha \in R$  we associate the corresponding reflection  $s_\alpha = s_{-\alpha} \in W$ . A root subsystem of  $R$  is by definition a subset  $R'$  of  $R$  stable under every  $s_\alpha, \alpha \in R'$ . The subgroup of  $W$  generated by the  $s_\alpha$  for  $\alpha \in R'$  is a reflection subgroup, and the map  $R' \mapsto \langle s_\alpha, \alpha \in R' \rangle$  defines a bijection between the set  $\mathcal{L}_R$  of all root subsystems and  $\mathcal{L}_\infty$ . The preordering induced by this bijection on  $\mathcal{L}_R$  is simply the inclusion ordering. We endow  $\mathcal{L}_R$  with the corresponding join semilattice structure. The cyclic reflection subgroups of  $W$  correspond to the root subsystems  $\{\alpha, -\alpha\}$  for  $\alpha \in R$ .

We let  $\mathcal{L}_c$  denote the subset of  $\mathcal{L}_R$  corresponding to the *closed* subsystems, namely the  $R' \in \mathcal{L}_\infty$  for which  $\forall \alpha, \beta \in R' \quad \alpha + \beta \in R \Rightarrow \alpha + \beta \in R'$ . Note that an intersection of closed subsystems is a closed subsystem, and that the subsystems of the form  $\{\alpha, -\alpha\}$  as well as the empty subsystem are closed. We have a map  $c : \mathcal{L}_R \rightarrow \mathcal{L}_c$  which associates to  $R' \in \mathcal{L}_R$  its *closure*, namely the intersection of all closed subsystems containing it. It is immediately checked that  $c$  is  $W$ -equivariant and a join semilattice morphism. From this it follows that we get an admissible map  $\mathcal{L}_\infty \simeq \mathcal{L}_R \rightarrow \mathcal{L}_c$ .

This proves the following.

**Proposition 4.4.** *Let  $R$  be a reduced root system and  $W$  the associated finite Coxeter group. Under the identification  $\mathcal{L}_\infty \simeq \mathcal{L}_R$ , the map  $c : \mathcal{L}_R \rightarrow \mathcal{L}_c$  induces a surjective morphism  $\mathcal{C}(W, \mathcal{L}_\infty) \rightarrow \mathcal{C}(W, \mathcal{L}_c)$ .*

This proposition proves that the algebra  $\mathcal{C}(W, \mathcal{L}_c)$  is isomorphic to the algebra  $\mathcal{C}_W^R$  of [18], which generically embeds into the corresponding Yokonuma-Hecke algebra. Indeed,  $\mathcal{C}_W^R$  is defined as a quotient of  $\mathcal{C}(W, \mathcal{L}_\infty) = \mathcal{C}_W$ , and one gets immediately that the map  $\mathcal{C}_W \rightarrow \mathcal{C}(W, \mathcal{L}_c)$  defined above factors through  $\mathcal{C}_W \rightarrow \mathcal{C}_W^{(R)}$ . The induced surjective map  $\mathcal{C}_W^{(R)} \rightarrow \mathcal{C}(W, \mathcal{L}_c)$  is then checked to be injective, since the natural spanning set of  $\mathcal{C}_W^{(R)}$  is mapped to a basis of  $\mathcal{C}(W, \mathcal{L}_c)$ . It is then immediately checked that the corresponding diagram of isomorphisms and natural projections is commutative.

$$\begin{array}{ccccc}
 \mathcal{C}_W & \xrightarrow{\simeq} & \mathcal{C}(W, \mathcal{L}_\infty) & & \\
 \downarrow & \searrow & \downarrow & \searrow & \\
 & & \mathcal{C}_W^{(R)} & \xrightarrow{\simeq} & \mathcal{C}(W, \mathcal{L}_c) \\
 \downarrow & \swarrow & \downarrow & \swarrow & \\
 \mathcal{C}_W^{(p)} & \xrightarrow{\simeq} & \mathcal{C}(W, \mathcal{L}_p) & & 
 \end{array}$$

**4.6. A priori unrelated examples.** A computer-aided exploration shows that there are other admissible lattices not originating a priori from root systems, with  $\mathcal{L}_p \subset \mathcal{L} \subset \mathcal{L}_\infty$ . In type  $A$  we have  $\mathcal{L}_p = \mathcal{L}_\infty$ , but in type  $D_n$  for  $n \geq 4$  we have  $\mathcal{L}_p \subsetneq \mathcal{L}_\infty$  while all root subsystems are closed. We checked for small  $n$  whether there are other admissible lattices in type  $D_n$ . This can be done as follows. First of all, one computes the  $W$ -orbits for the action on  $\mathcal{L}_\infty \setminus \mathcal{L}_p$ , since  $\mathcal{L}_\infty \setminus \mathcal{L}$  has to be a union of them. For each such union of orbits we then test whether the obtained subset  $\mathcal{L}$  satisfies the join semilattice property. In type  $D_4$ , the action of  $W$  on  $\mathcal{L}_\infty \setminus \mathcal{L}_p$  is transitive (and it is the orbit of a reflection subgroup of type  $A_1^4$ ), so there is no intermediate admissible lattice. But in type  $D_5$ , the action has 2 orbits, one of type  $A_1^4$  inherited from type  $D_4$ , and the other one of type  $A_1 A_3$ . By adding to  $\mathcal{L}_p$  the orbit of type  $D_4$  one checks by computer that the corresponding poset  $\mathcal{L}$  is admissible, every two elements admitting a join. This proves that examples containing the lattice of parabolic subgroups and which are a priori not related to the theory of root systems do exist.

## REFERENCES

- [1] F. Aicardi, J. Juyumaya, *An algebra involving braids and ties*, ICTP Preprint IC/2000/179.
- [2] F. Aicardi, J. Juyumaya, *Markov trace on the algebra of braids and ties*, Moscow Math. J. **16**(3) (2016) 397–431.
- [3] S. Ariki, *Representation theory of a Hecke algebra of  $G(r, p, n)$* , J. Algebra **177** (1995), 164–185.
- [4] S. Ariki, K. Koike, *A Hecke algebra of  $(\mathbf{Z}/r\mathbf{Z}) \wr \mathfrak{S}_n$  and construction of its irreducible representations*, Advances in Math. **106** (1994), 216–243.
- [5] D. Bessis, *Variations on Van Kampen’s method*, J. Math. Sci. (N.Y.) **128** (2005), 3142–3150.
- [6] E. Chavli, *The BMR freeness conjecture for exceptional groups of rank 2*, doctoral thesis, Univ. Paris Diderot (Paris 7), 2016.
- [7] E. Chavli, *The BMR freeness conjecture for the first two families of the exceptional groups of rank 2*, to appear in Comptes Rendus Mathématiques.
- [8] E. Chavli, *Universal deformations of the finite quotients of the braid group on 3 strands*, J. Algebra **459** (2016), 238–271.
- [9] M. Broué, G. Malle, J. Michel, *Towards Spetses I*, Transform. Groups **4** (1999), 157–218.
- [10] M. Broué, G. Malle, R. Rouquier, *Complex reflection groups, braid groups, Hecke algebras*, J. Reine Angew. Math. **500** (1998), 127–190.
- [11] M. Cabanes, M. Enguehard, *Representation theory of finite reductive groups*, Cambridge, 2004.
- [12] M. Cabanes, I. Marin, *On ternary quotients of cubic Hecke algebras*, Comm. Math. Phys. **314** (2012), 57–92.
- [13] J.I. Cogolludo-Agustín, *Braid monodromy of algebraic curves*, Ann. Math. Blaise Pascal **18** (2011), 141–209.
- [14] J. Espinoza, S. Ryom-Hansen, *Cell structures for the Yokonuma-Hecke algebra and the algebra of braids and ties*, preprint 2016, arxiv 1503.03396v3.
- [15] I. Marin, *The cubic Hecke algebra on at most 5 strands*, J. Pure Appl. Algebra **216** (2012), 2754–2782.
- [16] I. Marin, *Krammer representations for complex braid groups*, J. Algebra **371** (2012), 175–206.
- [17] I. Marin, *The freeness conjecture for Hecke algebras of complex reflection groups, and the case of the Hessian group  $G_{26}$* , J. Pure Appl. Algebra **218** (2014), 704–720.

- [18] I. Marin, *Artin groups and the Yokonuma-Hecke algebras*, preprint 2016, to appear in I.M.R.N.
- [19] I. Marin, *Proof of the BMR freeness conjecture for the groups  $G_{20}$  and  $G_{21}$* , preprint 2017, arXiv:1701.09017.
- [20] I. Marin, G. Pfeiffer, *The BMR freeness conjecture for the 2-reflection groups*, preprint 2014, to appear in Math. of Comput.
- [21] P. Orlik, H. Terao, *Arrangements of hyperplanes*, Springer-Verlag, Berlin, 1992.
- [22] S. Ryom-Hansen, *On the representation theory of an algebra of braids and ties*, J. Algebraic Combin. **33** (2011), no. 1, 57–79.
- [23] J.-P. Serre, *Linear representations of finite groups*, GTM **42**, Springer, 1977.
- [24] R. P. Stanley, *Enumerative combinatorics, Volume 1*, 2nd edition, Cambridge University Press, 2012.
- [25] T. Yokonuma, *Sur la structure des anneaux de Hecke d'un groupe de Chevalley fini*, C. R. Acad. Sci. Paris Série A-B **264** (1967), A344-A347.

LAMFA, UNIVERSITÉ DE PICARDIE-JULES VERNE, AMIENS, FRANCE  
*E-mail address:* `ivan.marin@u-picardie.fr`